

The 19th Congress of the European Association of Hospital Pharmacists, Barcelona, March 2014



# The art of writing an abstract

**Lene Juel Kjeldsen,** PhD Senior Researcher in Clinical Pharmacy The Research Unit for Hospital Pharmacy Amgros, Denmark





Conflict of interest: Nothing to disclose





# Pimp the abstract - agenda

Example 1 of rejected abstract

#### ►Group work

 Pimp the abstract to apply with the EAHP criteria for abstract content (Example 2 of rejected abstract)



2





# **Example 1 I**

#### **▶**Title

The use of alternative lipid emulsions in pediatric parenteral nutrition

## **▶**Background

 Lipid emulsions are an important part of pediatric parenteral nutrition, providing energy, fat soluble vitamins and they are a source of essential fatty acids. Soyabean oil emulsions have however been identified as one of the major causes of PN-associated cholestasis (PNAC) development, caused by peroxidation and free radical formation. Long-term PN is related to several complications as liver diseases, which might progress to cirrhosis and liver failure.

**CAMGROS** 



## **Example 1 II**

#### ▶Purpose

- Soyabean oil has been the standard lipid emulsion used in parenteral nutrition for many years without alternatives. The use of fish oil has been proposed as a possible alternative to soyabean oil, especially in treating and/or preventing PNAC. A newer lipid emulsion, based on fish oil, has been shown to reduce the risk of cholestasis but this practice isn't recommended because it could be detrimental to infants who have limited fat stores.



**CAMGROS** 



## **Example 1 III**

#### ▶ Materials and methods

- Soyabean oil is composed of long chain triglycerides and contains > 60% polyunsaturated fatty acids, with an omega-6:omega-3 ratio of 5.5:1. An addiction of fish oil instead provides a reduction of omega-6 fatty acids and a higher intake of omega-3 fatty acids, which possess potent immunomodulatory and anti-inflammatory properties. However, both omega-6 and omega-3 fatty acids have been shown to be essential for growth and development. Olive oil emulsions instead wouldn't provide sufficient omega-3 fatty acids.







**CAMGROS** 

.



## **Example 1 IV**

#### ▶ Results

 Recent studies have suggested that fish oil might lead to faster reversal of cholestasis than soyabean oil, reporting a median time of 9.4 weeks to reversal of cholestasis vs.
44.1 weeks of a control group receiving soyabean oil (4.69 times faster). Evidence from controlled trials have shown that fish oil combined with other lipids could provide similar beneficial effects.

#### **▶**Conclusions

 The use of multi-lipid emulsions is advised to prevent and/or treat PNAC, combining the beneficial aspects of different lipid emulsions as a safe and well tolerated vehicle.

6





## **Example 1 V**

### ▶ Potential improvements

- The title does not cohere with the data/conclusion of the study
- Background information is not limited to the background section
- The purpose is missing
- The materials and methods are not presented
- The type of study is unclear literature review?
- The results do not represent findings related to the purpose and methods (not) presented
- The conclusion does not strictly address the (non existing) purpose
- Lack of clear explanation of abbreviations
- What does the reader gain from the abstract?

**CAMGROS** 

7



# Pimp the abstract I

- ▶ Pimp the abstract to comply with the EAHP criteria for abstract content
  - What is the story presented in the abstract?
  - What can be improved?
  - How can the text be improved?
    - · Suggest new sentences
    - Focus on structure not too much on the facts presented





# Pimp the abstract II

#### **▶**Title

- DEVELOPMENT AND IMPLEMENTATION OF A PHARMACOTHERAPEUTIC PROTOCOL IN PATIENTS SUBMITTED TO CAESAREAN

## **▶**Background

 The medication errors are the most frecuent cause of adverse events. To develop pharmacotherapeutic protocols are a usefull tool in improving the quality and safety patient care.




**CAMGROS** 



# Pimp the abstract III

#### ▶ Materials and methods

- Prospective study of use of carbetocin in patients submitted to caesarean during a period of seven months. An interdisciplinary team was created and a chronogram of work was described. A bibliographic search was performed using Medline. The updated scientific evidence was evaluated. Other information sources used:product information, clinical guidelines. The protocol was established. The adequacy of the carbetocin use to the established conditions at protocol were analyzed.



10





## Pimp the abstract IV

#### ▶ Results

-Inclusion criteria, conditions of prescription and dispensing process were established. All carbetocin treatments were analyzed on the basis on information collected through the request form and the patient clinical records. 966 childbirths occured. 164(17%) were performed by caesarean. In 9(5,5%) carbetocin was used. In all cases, request form was perfectly filled out, checking previously if every case met the protocol. All carbetocin treatments were accord with protocol.

**CAMGROS** 



# Pimp the abstract V

#### ▶Conclusions

- The protocol of use of carbetocin in the hospital has been followed adequately according to the established guidelines. The use of this drug has been limited to patients in whom carbetocin treatment is the most efficient and safe option. Moreover, unnecessary risks and costs are avoided. The protocol let the hospital having a record and a follow-up of the patients, which makes possible the continuous assessment of the economic and clinical results. Results should be assesed for a minimum of one year in order to obtain more reliable data and to identify new opportunities of improvement.









## Pimp the abstract VI

### ▶ Potential improvements

- The title and purpose do not cohere with the conclusion of the study
- The background does not clearly describe the problem
- The purpose contains 2-3 aims
- The materials and methods do not clearly describe, which methods are used to address each of the aims
- The results contain inclusion criteria (should be in the methods section and related to the relevant aim)
- The results do not clearly indicate, which data are related to the relevant aims
- The conclusion does not address all aims
- The conclusion contains new information
- The conclusion is contradictive regarding the concluding statements and future research
- General: Grammar, spelling mistakes etc.





# Thank you for your time & attention



14

